



Short Note

Micronutrient cations status in vegetable growing soils of sub-humid and wet-temperate zones of Himachal Pradesh

**Girish Chander¹, Sandeep Sharma², Vivek Sharma³, Sudhir Verma⁴,
Sheetal Sharma⁵ and T.S. Verma**

Department of Soil Science
CSK Himachal Pradesh Krishi Vishvavidyalaya, Palampur-176 062, India.

Manuscript Received: 14.01.2014; Accepted: 02.07.2014

Abstract

Composite surface (0 to 0.15 m) soil samples were collected from vegetable growing fields in mid-hills sub-humid and high-hills wet-temperate sub-agroclimatic zones of Himachal Pradesh. The soils in wet temperate zone of Himachal Pradesh were more acidic (pH 5.2 to 6.8) in contrast to soils in mid-hills sub-humid zone (pH 5.8 to 7.5) sandy loam to clay loam in texture, medium to high in organic carbon content. Soils were medium in available-N, low to medium in available-P and medium to high in available-K. In sub-humid zone, the DTPA-extractable micronutrients ranged between 0.64 to 11.0 mg kg⁻¹ for Zn, 0.14 to 2.80 mg kg⁻¹ for Cu, 10.6 to 70.8 mg kg⁻¹ for Fe and 2.1 to 34.9 mg kg⁻¹ for Mn, while in wet-temperate zone between 0.44 to 2.06 mg kg⁻¹ for Zn, 0.02 to 3.60 mg kg⁻¹ for Cu, 22.8 to 96.6 mg kg⁻¹ for Fe and 2.5 to 40.0 mg kg⁻¹ for Mn. Soils in some parts of wet temperate zone of Himachal Pradesh are inherently low in DTPA-extractable Zn (19%), Cu (13%) and Mn (6%) while parts of sub humid zone are low in DTPA-extractable Cu (16%) and Mn (5%). All the DTPA micronutrients showed positive correlation with organic carbon. DTPA-Fe and Mn showed a negative correlation with soil pH. Among mechanical separates, the availability of DTPA-Cu and Mn was higher with finer soil fractions and decreased with coarser ones and reverse relation was observed with DTPA-Zn.

Key words: Micronutrients, DTPA- extractable Zn, DTPA- extractable Fe, DTPA- extractable Mn, DTPA-extractable Cu.

Micronutrients are essential for plant growth, yield maximization and quality improvement of crop produce (Kumar *et al.* 2004; Wani *et al.* 2003), but have not received the required attention like

macronutrients for sustainable crop intensification. Studies have indicated widespread deficiencies of micronutrients along with macronutrients limiting the realization of crop productivity potential (Wani *et.*

¹Resilient Dryland Systems, International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru-502324, Andhra Pradesh, India

^{2,3}Department of Soil Science, Punjab Agricultural University, Ludhiana 141 004, India

⁴Department of Soil Science and Water Management, Dr YS Parmar University of Horticulture and Forestry, Nauni, Solan, HP, India

⁵International Rice Research Institute (IRRI), India-Office, NASC, New Delhi, India

al. 2009, 2013; Sahrawat *et al.* 2007, 2010; Girish Chander *et al.* 2012, 1013). The reasons for occurrence of such deficiencies are nutrient mining under intensive farming and undue focus on high analysis NPK fertilizers while ignoring the micronutrients. Micronutrients are no doubt required in smaller quantities, but any deficiency acts as a weakest link inhibiting realization of productivity potential and sustainability. The inherent capacity of soils to supply micronutrients primarily governs their vulnerability for occurrence of such deficiencies and therefore needs desired attention to assess sensitive regions for resorting to appropriate fertilizer management strategies to apply deficient micronutrients from external sources. However, such information on soils in sub-humid and wet-temperate regions of Himachal Pradesh is very limited. Sub-humid region occupies 32% of the total geographical area (55,67,300 ha) and 37% of the net cultivated area (5,60,154 ha) of the state, while wet-temperate region covers 35% of the geographical area and 21% of the cultivated area of the state (DoAHP, 2011). Vast tracts in these regions have diversified into high input based seasonal and off-season vegetable production and are prone to setbacks in yield and sustainability due to heavy nutrient mining and deterioration in soil fertility. The present study was undertaken to determine the DTPA-extractable micronutrients status of undisturbed soils in sub-humid and wet-temperate regions of Himachal Pradesh.

Composite surface (0 to 0.15 m) soil samples were collected from vegetable growing fields in mid-hills sub-humid and high-hills wet-temperate sub-agroclimatic zones of Himachal Pradesh. The undisturbed soils were used for this purpose with a view to avoid the effects of different cultivation and fertilization practices. The soil samples so collected were air dried, ground with the help of wooden pestle and mortar, sieved through 2 mm sieve and analyzed for pH (Glass Electrode Method, Jackson, 1967), organic carbon (Walkley and Black rapid titration method, Walkley and Black, 1934), texture

(International Pipette Method, Black, 1965), Exchangeable calcium (Black, 1965), available nitrogen (Alkaline Permanganate Method, Subbiah and Asija, 1956), available phosphorus (Olsen *et al.* 1954) and available potassium (AOAC, 1970). The available Zn, Cu, Fe and Mn were estimated by the DTPA method (Lindsay and Norvell, 1978) on atomic absorption spectrophotometer. Simple correlation of DTPA-extractable micronutrients with these soil properties were worked out as per detailed procedure prescribed by Snedecor and Cochran (1994). The findings are discussed below:

Physico-chemical properties of soil

The soils in wet temperate zone of Himachal Pradesh were more acidic (pH 5.2 to 6.8) in comparison to soils in mid-hills sub-humid zone (pH 5.8 to 7.5) which might be due to more leaching of bases because of high precipitation. Soil texture was sandy loam to clay loam in both the zones; however sandy loam and loam were the dominated textured in majority of soils (Table 1). Soils were medium to high in organic carbon content due to luxurious plant growth and slow rate of organic matter decomposition due to low temperature. Majority of soils were medium in available-N, low to medium in available-P and medium to high in available-K. The soils of high-hills wet-temperate zone contained more available-N and lesser contents of available-P and K in comparison to soils in sub humid zone. However, the contents of exchangeable-Ca were also lower in high-hills wet-temperate zone soils in contrast to mid-hills sub-humid zone soils.

Micronutrient cations status in soil

In sub-humid zone, the DTPA-extractable micronutrients ranged between 0.64 to 11.0 mg kg⁻¹ for Zn, 0.14 to 2.80 mg kg⁻¹ for Cu, 10.6 to 70.8 mg kg⁻¹ for Fe and 2.1 to 34.9 mg kg⁻¹ for Mn, while in wet-temperate zone between 0.44 to 2.06 mg kg⁻¹ for Zn, 0.02 to 3.60 mg kg⁻¹ for Cu, 22.8 to 96.6 mg kg⁻¹ for Fe and 2.5 to 40.0 mg kg⁻¹ for Mn (Table 2). The results showed in general higher contents of DTPA-Fe and Mn in wet-temperate region soils as comparison to

Table 1. Soils physico-chemical properties in mid-hills sub-humid and high-hills wet-temperate zones of Himachal Pradesh

Property	Mid hills sub-humid zone	High-hills wet-temperate zone
pH	5.8-7.5	5.2-6.8
Oxid. Org. C (g kg ⁻¹)	9.3-27.5	9.5-22.7
Exch. Ca (meq/kg)	276-2195	761-1611
Clay (%)	8.9-38.6	10.0-34.8
Silt (%)	12.3-46.5	10.7-41.6
Sand (%)	22.4-78.8	12.1-77.0
Textural Class	Sandy loam – clay loam	Sandy loam – clay loam
Available nitrogen (kg ha ⁻¹)	282-439	282-502
Available phosphorus (kg ha ⁻¹)	2.3-37.0	2.3-37.0
Available potassium (kg ha ⁻¹)	39-560	67-504

Table 2. DTPA-extractable micronutrient status (mg kg⁻¹) of soils in mid-hills sub-humid and high-hills wet-temperate zones of Himachal Pradesh

	Mid-hills sub-humid zone				High-hills wet-temperate zone			
	Zn	Cu	Fe	Mn	Zn	Cu	Fe	Mn
Range	0.64 - 11.0	0.14 - 2.80	10.6 – 70.8	2.1 – 34.9	0.44 – 2.06	0.02 – 3.60	2.8 – 96.6	2.5 – 40.0
Mean	3.36	1.07	35.3	17.5	0.94	1.01	47.5	21.1
% deficiency	0	16	0	5	19	13	0	6

Table 3. Relationship between soil properties and DTPA- extractable micronutrients

Soil property	Correlation coefficient (r)			
	Zn	Cu	Fe	Mn
Clay	-0.14	0.14	-0.01	0.18
Silt	-0.28	-0.10	0.08	0.33
Sand	0.27	-0.01	-0.03	-0.29
Organic-C	0.04	0.12	0.13	0.01
pH	0.18	0.16	-0.66**	-0.22

that in sub-humid zone which might be because of more acidic nature of the wet-temperate soils (Average soil pH was 6.1 in wet-temperate zone and 6.5 in sub-humid zone) as they have developed under the influence of high rainfall and coniferous type of vegetation. Considering the critical limits of 0.5 mg kg⁻¹ for DTPA-Zn, 0.2 mg kg⁻¹ for DTPA-Cu, 4.5 mg kg⁻¹ for DTPA-Fe (Follet and Lindsay, 1970) and 3.5 mg kg⁻¹ for DTPA-Mn, the soils were sufficient in DTPA-Zn and Fe under sub-humid zone and in DTPA-Fe under wet-temperate zone. In sub-humid

zone, about 16 per cent soils had low level of DTPA-Cu and 5 per cent had low DTPA-Mn levels confined in Kangra and Solan districts. The Solan district was critical for both DTPA-Cu and Mn, while Kangra for DTPA-Cu only. Similarly, in wet-temperate zone the low levels of DTPA micronutrients were recorded for Zn (19 per cent), Cu (13 per cent) and Mn (6 per cent) in the districts of Shimla and Kullu. The Shimla district witnessed low levels of DTPA-Zn and Cu, while Kullu district that of DTPA-Zn and Mn. In the regions with detected low levels of micronutrients,

intensive vegetable cultivation is being done which is apparently leading to heavy nutrient mining and raises questions over the sustainability and profitability.

Relationship of soil micronutrient cations with soil properties

In an endeavor to understand the relationship with soil properties, an attempt was made to correlate DTPA-extractable micronutrients in sub-humid and wet-temperate zones with soil mechanical separates, organic-C and pH (Table 3). All the DTPA micronutrients showed positive correlation with organic C suggesting that their availability is high and is controlled by this soil component. DTPA-Fe and Mn showed a negative correlation with soil pH indicating that their availability is largely influenced by acidic nature of the soil. But a significant relationship was observed between DTPA-Fe and pH ($r = -0.66$) only. A similar relationship between DTPA-Fe and pH has also been reported by Kanwar *et al.* (1986) and Sharma *et al.* (2003). Reduction in availability of Fe with an increase in pH may be attributed to precipitation as insoluble $\text{Fe}(\text{OH})_2$. As regards the relationship with mechanical separates (sand, silt & clay), the availability of DTPA-Mn was higher with finer soil fractions and decreased with

coarser ones. Finer soil fractions provide sites on which plant nutrients are adsorbed and thus have more nutrient holding capacity as compared to coarse textured fractions. As regards the relationship between mechanical separates and DTPA-Zn, a reverse relation was observed where the availability was less with finer soil fraction. This may be due to the oxides of Mn and Fe which have a significant effect on micronutrients particularly Zn reactions in soil resulting in irreversible retention of Zn (Das, 2000). However, relationships between soil properties and micronutrients except that for pH and DTPA-Fe were not statistically significant (Table 3). Soils in some parts of wet temperate zone of Himachal Pradesh are inherently low in DTPA-extractable Zn (19 per cent), Cu (13 per cent) and Mn (6 per cent) while parts of sub humid zone are low in DTPA-extractable Cu (16 per cent) and Mn (5 per cent). These areas are hub of intensive vegetable cultivation and therefore need extensive delineation of on-farm deficiencies and designing fertilizer management strategies that take into account emerging deficiencies to get higher productivity and quality on sustainable basis without degrading natural resources.

References

- AOAC 1970. *Official Method of Analysis of the Association of Analytical Chemists*. Benjamin Franklin, Station, Washington, DC. (USA).
- Black CA 1965. *Methods of Soil Analysis*, Part – I. American Society of Agronomy, Madison Wisconsin, USA, pp 900-01.
- Das DK 2000. *Micronutrients: Their Behavior in Soils and Plants*. Kalyani Publishers, New Delhi, pp 47-126.
- DoAHP (Department of Agriculture, Himachal Pradesh) 2011. *Agro climatic zones*. <http://www.hpagriculture.com/agro.htm>.
- Follet RH and Lindsay WL 1970. Profile distribution of zinc, iron, manganese and copper in Colorado soils. Colorado State University Experimental Station Tech. Bull. **110**: 79.
- Girish Chander, Wani SP, Sahrawat KL and Jangawad LS 2012. Balanced plant nutrition enhances rainfed crop yields and water productivity in Jharkhand and Madhya Pradesh states in India. *J. Trop. Agric.* 50 (1-2): 24-29.
- Girish Chander, Wani SP, Sahrawat KL, Kamdi PJ, Pal CK, Pal DK and Mathur TP 2013. Balanced and integrated nutrient management for enhanced and economic food production: case study from rainfed semi-arid tropics in India. *Arch. Agron. Soil Sci.* doi: <http://dx.doi.org/10.1080/03650340.2012.761336>.
- Jackson ML 1967. *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd. New Delhi.
- Kanwar BB, Tripathi BR and Kanwar BS 1986. Distribution of total and DTPA-extractable iron in

- relation to soil properties in some soil profiles of North-Western Himalayas. *Him. J. Agric. Res.* **12**: 108-18.
- Kumar PSS, Geetha SA, Savithri P and Mahendran PP 2004. Influence of different micronutrient treatments on nutrient status of the soil, yield, growth and quality parameters in Muscat grapes (*Vitis vinifera* var. L.). *Res. Crops* **5**: 85-95.
- Lindsay WL and Norvell WA 1978. Development of a DTPA soil test for zinc, iron, manganese, and copper. *Soil Sci. Soc. Am. J.* **42**: 421-28.
- Olsen SR, Cole VC, Watanabe FS and Dean LA 1954. Estimation of available phosphorus in soil by extraction with NaHCO₃, USDA and Circ. **3**: 19-33.
- Sahrawat KL, Wani SP, Rego TJ, Pardhasaradhi G and Murthy KVS 2007. Widespread deficiencies of sulphur, boron and zinc in dryland soils of the Indian semi-arid tropics. *Current Sci.* **93**: 1428-32.
- Sahrawat KL, Wani SP, Parthasaradhi G and Murthy KVS 2010. Diagnosis of secondary and micronutrient deficiencies and their management in rainfed agroecosystems: case study from Indian semi-arid tropics. *Comm. Soil Sci. Plant Anal.* **41**: 346-60.
- Sharma RP, Megh Singh and Sharma JP 2003. Correlation studies on micronutrients vis-à-vis soil properties in some soils of Nagaur district in semi arid region of Rajasthan. *J. Indian Soc. Soil Sci.* **51**: 522-27.
- Snedecor GW and Cochran WG 1994. *Statistical Methods*, Eighth edition, Affiliated East-West Press Pvt. Limited, New Delhi, pp 503.
- Subbiah BV and Asija GL 1956. A rapid procedure for estimation of available nitrogen in soils. *Current Sci.* **25**: 259-60.
- Walkley A and Black IA 1934. An examination of the Degtjareff method for determination of soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci.* **37**: 29-38.
- Wani SP, Singh P, Boomiraj K and Sahrawat KL 2009. Climate change and sustainable rain-fed agriculture: challenges and opportunities. *Agric. Situation in India* **66**:221-39.
- Wani SP, Girish Chander, Sahrawat KL, Dixit S and Venkateswarlu B 2013. Improved crop productivity and rural livelihoods through balanced nutrition in the rain-fed semi-arid tropics. *Resilient Dryland Systems Report No. 58*. Patancheru, Andhra Pradesh.
- Wani SP, Pathak P Jangawad LS, Eswaran H and Singh P 2003. Improved management of vertisols in the semiarid tropics for increased productivity and soil carbon sequestration. *Soil Use and Management* **19**:217-22.